

Z' and anomalous gauge coupling effects at LEP2 and their separation

A. A. Babich¹

International Centre for Theoretical Physics, Trieste, Italy

A. A. Pankov²

International Centre for Theoretical Physics, Trieste, Italy
Istituto Nazionale di Fisica Nucleare, Sezione di Trieste, Trieste, Italy

N. Paver³

Dipartimento di Fisica Teorica, Università di Trieste, Trieste, Italy
Istituto Nazionale di Fisica Nucleare, Sezione di Trieste, Trieste, Italy

Abstract

In the case of the process $e^+e^- \rightarrow \mu^+\mu^-$, we study the sensitivity to either Z' or anomalous gauge coupling effects of two new observables, σ_+ and σ_- , conveniently defined for such kind of analysis. We discuss general properties of the deviations of these observables from the Standard Model predictions, and derive the model-independent bounds on the relevant parameters allowed by present and foreseeable LEP2 data. We also discuss the possibility of separating the two kinds of effect.

¹Permanent address: Department of Mathematics, Polytechnical Institute, Gomel, 246746 Belarus. E-mail: BABICH@GPI.GOMEL.BY

²Permanent address: Department of Physics, Polytechnical Institute, Gomel, 246746 Belarus. E-mail: PANKOV@GPI.GOMEL.BY

³Also supported by the Italian Ministry of University, Scientific Research and Technology (MURST).

1 Introduction

The existence of a heavy neutral gauge boson Z' is theoretically motivated by most extended electroweak scenarios attempting to overcome the conceptual difficulties of the Standard Model [1]. The knowledge of the Z' parameters, such as the mass and couplings to ordinary fermions, would be essential in order to test these extended theories. Numerous strategies to evidence manifestations of the Z' in experiments at high energy e^+e^- and hadronic colliders have been developed. Due to the present ‘direct’ lower mass limit from the Tevatron, $M_{Z'} > \mathcal{O}(500)$ GeV [2], only ‘indirect’ manifestations of the Z' can be searched for at the energy of LEP2. The preferred channel in this case should be the electron-positron annihilation into fermion pairs, where the Z' contributes already at the Born level [3]. In this regard, particularly convenient should be the annihilation into lepton pairs

$$e^+ + e^- \rightarrow \bar{l} + l \quad (l = \mu, \tau) \quad (1)$$

where, assuming lepton universality, the number of Z' parameters is significantly reduced, leading to characteristic, model-independent potential signals. These effects manifest themselves as deviations of observables from the Standard Model (SM) predictions, to be compared to the experimental data, not only as regards the size of the signals, but also their characteristic signs and energy dependences. In the case of no observed signal within the experimental accuracy, the result of the analysis can be expressed as bounds on the Z' parameters to a conventionally specified confidence level.

Similar effects in process (1) can arise from the independent physical mechanism represented by anomalous gauge boson couplings, that can contribute even at the tree level [4]. Clearly, in the case of no observed effect, upper limits on the anomalous gauge couplings can be obtained from the data. On the other hand, it would be desirable to have some criterion to distinguish the two sources of such nonstandard effects, in the case these were observed.

The aim of this paper is to present a discussion of the general properties of the deviations from the SM predictions due to the Z' and to the anomalous gauge couplings, at the energies of LEP2. In particular, some criterion to distinguish the two kinds of signal is worked out. Moreover, the constraints on the relevant parameters are derived either from the present LEP2 data and in the case of the foreseeable ones.

2 Manifestations of Z'

We start from the general neutral-current interaction including the Z' :

$$-L_{NC} = eJ_\gamma^\mu A_\mu + g_Z J_Z^\mu Z_\mu + g_{Z'} J_{Z'}^\mu Z'_\mu, \quad (2)$$

where $e = \sqrt{4\pi\alpha_{e.m.}}$; $g_Z = e/s_W c_W$ (with $s_W^2 = 1 - c_W^2 \equiv \sin^2 \theta_W$) and $g_{Z'}$ are the Z and Z' gauge couplings, respectively. The currents can be written as

$$J_i^\mu = \sum_f \bar{\psi}_f \gamma^\mu (L_i^f P_L + R_i^f P_R) \psi_f = \sum_f \bar{\psi}_f \gamma^\mu (V_i^f - A_i^f \gamma_5) \psi_f. \quad (3)$$

Here, $i = \gamma, Z, Z'$ and $P_{L,R} = (1 \mp \gamma_5)/2$ are projectors onto the left- and right-handed electron helicity states, so that $R_i^f = V_i^f - A_i^f$ and $L_i^f = V_i^f + A_i^f$. The Standard Model (SM) couplings are

$$V_\gamma^f = Q_f; \quad A_\gamma^f = 0; \quad V_Z^f = \frac{I_{3L}^f}{2} - Q_f s_W^2; \quad A_Z^f = \frac{I_{3L}^f}{2}. \quad (4)$$

In general, the unpolarized differential cross section for electron-positron annihilation into a fermion-antifermion pair, $e^+e^- \rightarrow \bar{f}f$, can be written in Born approximation including γ, Z and Z' exchanges, as:

$$\frac{d\sigma_{ff}}{d\cos\theta} = \frac{\pi\alpha_{e.m.}^2}{2s} \left[(1 + \cos^2\theta) F_1 + 2\cos\theta F_2 \right], \quad (5)$$

where θ is the angle between the initial electron and the outgoing fermion.

For our purposes, it is convenient to split F_1 and F_2 as:

$$F_1 = F_1^{SM} + \Delta F_1, \quad F_2 = F_2^{SM} + \Delta F_2. \quad (6)$$

In Eq. (6):

$$\begin{aligned} F_1^{SM} &= Q_e^2 Q_f^2 + 2 Q_e v_e Q_f v_f \text{Re}\chi_Z + (v_e^2 + a_e^2)(v_f^2 + a_f^2)|\chi_Z|^2, \\ F_2^{SM} &= 2 Q_e a_e Q_f a_f \text{Re}\chi_Z + 4 v_e a_e v_f a_f |\chi_Z|^2, \end{aligned} \quad (7)$$

and the deviation due to the Z' is

$$\begin{aligned} \Delta F_1 &= 2 Q_e v'_e Q_f v'_f \text{Re}\chi_{Z'} + (v_e'^2 + a_e'^2)(v_f'^2 + a_f'^2)|\chi_{Z'}|^2 \\ &\quad + 2 (v_e v'_e + a_e a'_e)(v_f v'_f + a_f a'_f) \text{Re}(\chi_Z \chi_{Z'}^*), \\ \Delta F_2 &= 2 Q_e a'_e Q_f a'_f \text{Re}\chi_{Z'} + 4 v'_e a'_e v'_f a'_f |\chi_{Z'}|^2 \\ &\quad + 2(v_e a'_e + v'_e a_e)(v_f a'_f + v'_f a_f) \text{Re}(\chi_Z \chi_{Z'}^*). \end{aligned}$$

Here, the gauge boson propagators are $\chi_V = s/(s - M_V^2 + iM_V\Gamma_V)$, $V = Z, Z'$, and the fermionic coupling constants are normalized as:

$$v_f = \frac{g_Z}{e} V_Z^f, \quad a_f = \frac{g_Z}{e} A_Z^f, \quad v'_f = \frac{g_{Z'}}{e} V_{Z'}^f, \quad a'_f = \frac{g_{Z'}}{e} A_{Z'}^f. \quad (8)$$

The total width $\Gamma_{Z'}$ can be assessed from a sum over the partial widths:

$$\Gamma_{Z'}^{ff} = \frac{\alpha_{e.m.} M_{Z'}}{3} \sqrt{1 - 4m_f^2/M_{Z'}^2} \left[v_f'^2 + a_f'^2 + 2m_f^2/M_{Z'}^2 (v_f'^2 - 2a_f'^2) \right]. \quad (9)$$

The leptonic channel (1) has the nice phenomenological feature that, assuming leptonic $e-l$ universality, only three independent nonstandard parameters are needed: the couplings v'_l, a'_l and the mass $M_{Z'}$. More parameters would be needed in the description of $e^+e^- \rightarrow \bar{q}q$.

Concentrating on process (1) at LEP2 energies with $l = \mu$, the two conventionally used observables are the total cross section

$$\sigma_{\mu\mu} = \int_{-1}^1 \frac{d\sigma_{\mu\mu}}{d\cos\theta} d\cos\theta = \sigma_{\text{pt}} F_1, \quad (10)$$

with $\sigma_{\text{pt}} \equiv \sigma(e^+e^- \rightarrow \gamma^* \rightarrow \mu^+\mu^-) = (4\pi\alpha_{e.m.}^2)/(3s)$, and the forward-backward asymmetry

$$A_{\text{FB}} = \frac{\sigma_{\text{FB}}}{\sigma_{\mu\mu}} \equiv \frac{\sigma_{\mu\mu}^{\text{F}} - \sigma_{\mu\mu}^{\text{B}}}{\sigma_{\mu\mu}} = 3F_2/4F_1, \quad (11)$$

where the forward and backward cross sections are defined by $\sigma_{\mu\mu}^{\text{F}} = \int_0^1 (d\sigma_{\mu\mu}/d\cos\theta) d\cos\theta$ and $\sigma_{\mu\mu}^{\text{B}} = \int_{-1}^0 (d\sigma_{\mu\mu}/d\cos\theta) d\cos\theta$, respectively.

As it will appear in the sequel, in order to identify Z' effects in process (1) it may be convenient to consider, alternatively to A_{FB} , just the difference σ_{FB} between the forward and backward cross sections.

To discuss the distinctive features of nonstandard physical effects induced either by the exchange of a Z' or by anomalous gauge couplings, we introduce also the new observables σ_+ and σ_- defined as the differences of cross sections integrated in suitable ranges of $\cos\theta$:

$$\sigma_+ \equiv \left(\int_{-z^*}^1 - \int_{-1}^{-z^*} \right) \frac{d\sigma_{\mu\mu}}{d\cos\theta} d\cos\theta, \quad (12)$$

$$\sigma_- \equiv \left(\int_{-1}^{z^*} - \int_{z^*}^1 \right) \frac{d\sigma_{\mu\mu}}{d\cos\theta} d\cos\theta, \quad (13)$$

where $z^* > 0$ is determined from the condition that the kinematical coefficients multiplying F_1 and F_2 in Eq. (5) be the same after integration over $\cos\theta$. The condition is

$$\int_{-z^*}^{z^*} (1 + \cos^2\theta) d\cos\theta = \left(\int_{z^*}^1 - \int_{-1}^{-z^*} \right) 2\cos\theta d\cos\theta, \quad (14)$$

and gives the equation for z^*

$$\frac{1}{3}z^{*3} + z^{*2} + z^* - 1 = 0, \quad (15)$$

whose solution is $z^* = 2^{2/3} - 1 = 0.5874$, corresponding to $\theta^* = 54^\circ$. In the case of a reduced kinematical range of the scattering angle such as, e.g., $|\cos\theta| < c$, one derives $z^* = (1 + 3c)^{1/3} - 1$. One can write σ_+ and σ_- in terms of F_1 and F_2 as

$$\sigma_{\pm} = \sigma_{\text{pt}}^* (F_1 \pm F_2), \quad (16)$$

where, to a very good precision:

$$\sigma_{\text{pt}}^* \equiv \frac{3}{4} (1 - z^{*2}) \sigma_{\text{pt}} = 0.5 \sigma_{\text{pt}}. \quad (17)$$

The new, independent, observables $\sigma_{\pm}$ are simple combinations of the conventional ones $\sigma_{\mu\mu}$ and A_{FB} . Indeed, from Eqs. (10), (11), (16) and (17):

$$\sigma_{\pm} = 0.5 \sigma_{\mu\mu} \left(1 \pm \frac{4}{3} A_{\text{FB}} \right) = 0.5 \left(\sigma_{\mu\mu} \pm \frac{4}{3} \sigma_{\text{FB}} \right), \quad (18)$$

so that σ_+ and σ_- can be measured either directly according to Eqs. (12) and (13) or indirectly by means of $\sigma_{\mu\mu}$ and A_{FB} .

The previous formulae for the differential cross section (5), as well as those for all of the other observables, are still valid to a very good (improved Born) approximation which includes one-loop electro-weak radiative corrections, with the following replacements [5], [6]:

$$\begin{aligned} \alpha_{e.m.} &\Rightarrow \alpha(M_Z^2); & v_l &\Rightarrow \frac{1}{\sqrt{\kappa}} \left(I_{3L}^l - 2Q_e \sin^2 \theta_W^{\text{eff}} \right); & a_l &\Rightarrow \frac{I_{3L}^l}{\sqrt{\kappa}} \\ \sin^2 \theta_W &\Rightarrow \sin^2 \theta_W^{\text{eff}}; & \sin^2(2\theta_W^{\text{eff}}) &\equiv \kappa = \frac{4\pi\alpha(M_Z^2)}{\sqrt{2}G_F M_Z^2 \rho}; & \rho &\approx 1 + \frac{3G_F m_t^2}{8\pi^2 \sqrt{2}}, \end{aligned} \quad (19)$$

where only the main contribution to ρ , coming from the *top* mass, has been taken into account. This parameterization uses the best known SM parameters G_F , M_Z , and $\alpha(M_Z^2)$. Moreover, the energy dependence of the Z -width should be accounted for by the replacement of the Z propagator:

$$\chi_Z(s) \Rightarrow \frac{s}{s - M_Z^2 + i(s/M_Z^2)M_Z\Gamma_Z}. \quad (20)$$

In our analysis, the improved Born approximation with $m_t = 175$ GeV and $m_H = 300$ GeV is used.

While the numerical results are obtained from the exact formulae, for the sake of a simplified presentation of the analysis it is convenient to make a few, harmless, assumptions:

- i)* since the typical lower bound for the Z' boson mass, is much larger than the LEP2 energy, it suffices to take into account Z' interference effects only, the pure Z' exchange contributions being negligible;
- ii)* since in the SM $|v_l| \ll |a_l| < 1$, in the expressions of the deviations from the SM due to the Z' we neglect v_l with respect to a_l . In addition, one can neglect the imaginary part of the Z' boson propagator.

In the approximations *i)* and *ii)*, the deviation of the cross section from the SM prediction, $\Delta\sigma_{\mu\mu} = \sigma_{\mu\mu} - \sigma_{\mu\mu}^{SM}$, reads

$$\Delta\sigma_{\mu\mu} \approx 2\sigma_{\text{pt}} \left(v_l'^2 + a_l'^2 a_l^2 \text{Re } \chi_Z \right) \chi_{Z'}, \quad (21)$$

where the two terms represent the $\gamma - Z'$ and $Z - Z'$ interference, respectively, and v_l' , a_l' are the leptonic vector and axial-vector couplings to the Z' defined in Eq. (8).

In Eq. (21), only squares of leptonic couplings appear, so that one has the general property that the sign of the deviations of the cross section from the SM prediction

due to the Z' is definitely negative at the LEP2 energy, because both terms within the round brackets are positive. From Eq. (8), it is clear that this is independent of the simplifying assumption *ii*) above. Also, such sign definiteness holds only for process (1), assuming universality between initial and final leptons μ or τ , and not for the process $e^+e^- \rightarrow q\bar{q}$, where additional assumptions on the fermion coupling constants would be needed.

The deviation of σ_{FB} in (11) from the SM prediction, $\Delta\sigma_{\text{FB}} = \sigma_{\text{FB}} - \sigma_{\text{FB}}^{\text{SM}}$, has a similar expression:

$$\Delta\sigma_{\text{FB}} \approx \frac{3}{2} \sigma_{\text{pt}} \left(a_l'^2 + v_l'^2 a_l^2 \text{Re} \chi_Z \right) \chi_{Z'}, \quad (22)$$

and at LEP2 it generally has the same negative sign property for arbitrary leptonic Z' couplings as the cross section $\sigma_{\mu\mu}$. Such sign property of Z' effects for these observables can be used to attempt a model-independent distinction of the Z' from analogous anomalous gauge coupling effects, if some deviations were observed.

Concerning the forward-backward asymmetry (11), the deviation from the SM prediction is expressed as

$$\Delta A_{\text{FB}} = A_{\text{FB}} - A_{\text{FB}}^{\text{SM}} \propto \left[\left(a_l'^2 - \frac{4}{3} A_{\text{FB}}^{\text{SM}} v_l'^2 \right) + \left(v_l'^2 - \frac{4}{3} A_{\text{FB}}^{\text{SM}} a_l'^2 \right) a_l^2 \text{Re} \chi_Z \right] \chi_{Z'} \quad (23)$$

and, in contrast to the case of $\sigma_{\mu\mu}$ and σ_{FB} , at LEP2 energy it can be either positive or negative, depending on the actual values of the Z' couplings.

The deviation of σ_+ from the SM prediction has the expression

$$\Delta\sigma_+ = \sigma_+ - \sigma_+^{\text{SM}} \approx \sigma_{\text{pt}} (1 + a_l^2 \text{Re} \chi_Z) \cdot (v_l'^2 + a_l'^2) \chi_{Z'}. \quad (24)$$

Eq. (24) shows that the dependence of $\Delta\sigma_+$ on the Z' parameters is characterized by the expression $(v_l'^2 + a_l'^2) \chi_{Z'}$, which is a negative quantity at $\sqrt{s} < M_{Z'}$ for arbitrary Z' couplings. In addition to the definite negative sign, $\Delta\sigma_+$ has the property of depending on the total center of mass energy $\sqrt{s}$ via the factor $(1 + a_l^2 \text{Re} \chi_Z)$, which is completely determined by well-known SM parameters. Fig. 1a shows the energy dependence of the relative deviation $\Delta\sigma_+/\sigma_+^{\text{SM}}$. In this numerical example, we have considered a ‘sequential’ Z' [7] with couplings $v_l' = v_l$, $a_l' = a_l$ and with masses $M_{Z'} = 500$ GeV and $M_{Z''} = 700$ GeV. In this figure, one should notice the characteristic sign correlations of $\Delta\sigma_+$ at the different energies expected from a Z' ($\Delta\sigma_+/\sigma_+^{\text{SM}}$ is sequentially negative, positive and then negative again, with increasing $\sqrt{s}$).

As regards σ_- , it has in common with σ_+ the property that the deviation from the SM prediction

$$\Delta\sigma_- = \sigma_- - \sigma_-^{\text{SM}} \approx \sigma_{\text{pt}} (1 - a_l^2 \text{Re} \chi_Z) \cdot (v_l'^2 - a_l'^2) \chi_{Z'} \quad (25)$$

has an energy dependence completely determined in terms of SM parameters. However, in contrast to $\Delta\sigma_+$, the sign and magnitude of $\Delta\sigma_-$ is determined by $(v_l'^2 - a_l'^2) \chi_{Z'}$, for which both the positive and the negative signs are generally possible. Fig. 1b shows the energy dependence of the relative deviation $\Delta\sigma_-/\sigma_-^{\text{SM}}$, which,

Observables	Sign defined	Energy dependence defined
$\Delta\sigma_{\mu\mu}$	yes	no
$\Delta\sigma_{\text{FB}}$	yes	no
ΔA_{FB}	no	no
$\Delta\sigma_+$	yes	yes
$\Delta\sigma_-$	no	yes

Table 1: Model-independent properties of deviations induced by a Z' at LEP2 energy

similar to $\Delta\sigma_+$, has the characteristic sign correlations at the different energies. In the case of Fig. 1b, we take a Z' with same mass values as in Fig. 1a, but with couplings such that $v_l'^2 - a_l'^2 = \pm(v_l^2 - a_l^2)$ (to give an example).

The properties of the deviations $\Delta\mathcal{O}$ of the observables considered above ($\mathcal{O} = \sigma_{\mu\mu}, \sigma_{\text{FB}}, A_{\text{FB}}, \sigma_{\pm}$), are summarized in Table 1. As previously emphasized, these properties of the observables may be useful to identify (or to discard) the Z' as the (indirect) source of some deviations from the SM predictions, if experimentally observed.

Alternatively, it should be interesting to assess the sensitivity to the Z' of the observables $\mathcal{O}$, and the corresponding potential to constrain Z' parameters within an allowed region, in the case no deviations were observed within the expected experimental accuracy. We may attempt a model-independent analysis, which accounts either for the uncertainty of experimental data currently available from LEP2, or for the accuracy foreseeable for the future data with the planned integrated luminosity of 500 pb^{-1} . To this purpose, it is convenient to adopt the general expression of the neutral current interaction of the Z' given, e.g., in Refs. [8, 9], and rewrite (8) in terms of the ‘effective’ vector and axial-vector couplings

$$V_f = V_{Z'}^f \sqrt{\frac{g_{Z'}^2}{4\pi} \frac{M_Z^2}{M_{Z'}^2 - s}}, \quad A_f = A_{Z'}^f \sqrt{\frac{g_{Z'}^2}{4\pi} \frac{M_Z^2}{M_{Z'}^2 - s}}. \quad (26)$$

One of the advantages is that (26) allows to represent bounds on a two-dimensional scatter plot, without making reference to specific values of $M_{Z'}$ or s .

The sensitivity of the observables $\sigma_{\pm}$ has been assessed numerically by defining a χ^2 function as follows:

$$\chi^2 = \left(\frac{\Delta\mathcal{O}}{\delta\mathcal{O}} \right)^2, \quad (27)$$

where the uncertainty $\delta\mathcal{O}$ combines both statistical and systematic errors. As a criterion to derive allowed regions for the coupling constants in the case where no deviations from the SM were observed, and in this way to assess the sensitivity of process (1) to V_l and A_l through the observables $\mathcal{O}$, we impose that $\chi^2 < \chi_{\text{crit}}^2$, where χ_{crit}^2 is a number that specifies the desired ‘confidence’ level. We take the current integrated luminosities and systematic uncertainty relevant to the cross section of the leptonic process $e^+e^- \rightarrow \mu^+\mu^-$ at LEP2, as summarized in Table 2. In addition, to

$\sqrt{s}$	161 [GeV]	δ_{syst}	172 [GeV]	δ_{syst}
L3 [10]	10.9 pb^{-1}	4.0%	10.2 pb^{-1}	4.0%
OPAL [11]	10.1 pb^{-1}	2.2%	10.3 pb^{-1}	2.8%
ALEPH [12]	11.08 pb^{-1}	2.3%	10.65 pb^{-1}	2.3%
DELPHI [13]	9.95 pb^{-1}	3.5%	9.98 pb^{-1}	3.4%

Table 2: Systematic errors and integrated luminosities collected at current experiments at LEP2

assess future possibilities of improving the sensitivity, we work out the same example by using the expected final luminosity at LEP2, assuming for the systematic uncertainty, in this case, $\delta_{\text{syst}} = 0.5\%$ [3]). The numerical analysis has been performed by means of the program ZEFIT, which has to be used along with ZFITTER [14].

In Fig. 2 we show the results from $\sigma_{\pm}$, and compare the bounds on the couplings (26) obtainable from current and future experiments at LEP2, with inputs from Table 2. The contours are derived from the combination of both observables, σ_+ and σ_- , and correspond to two standard deviations ($\chi^2_{\text{crit}} = 4$). From the numerical point of view, a comparison of allowed bounds on (A_l, V_l) depicted in Fig. 2 with those obtained from a similar analysis of $\sigma_{\mu\mu}$ and A_{FB} indicates that the new observables σ_+ and σ_- have a slightly better sensitivity. This reflects the fact that, according to (24), (25) and (26), the deviations $\Delta\sigma_+$ and $\Delta\sigma_-$ are proportional to combinations of Z' coupling constants ($V_l^2 \pm A_l^2$, respectively) rather than the individual ones, so that a one-parameter fit applies in this case, rather than a two-parameter one. Moreover, we may remark that the above analysis is a model-independent one, as it starts from rather general considerations. Therefore, it can usefully complement the conventional analysis of Z' couplings based on the observables $\sigma_{\mu\mu}$ and A_{FB} .

3 Effects of the anomalous gauge couplings

In this regard, we now discuss a specific case of models with anomalous gauge couplings. This case was considered in Ref. [4], specifically concentrating on the $SU(2) \times U(1)$ invariant Lagrangian with only dimension six operators. Of these operators, four ones (denoted as $\mathcal{O}_{BW}$, $\mathcal{O}_{DW}$, $\mathcal{O}_{DB}$, and $\mathcal{O}_{\Phi,1}$) affect the neutral current and charged current amplitudes at tree level, and five more induce effects at the one-loop level. Of these five operators, three ones lead to anomalous trilinear vector boson couplings. As it was shown in Ref. [4], the quadratically and logarithmically divergent contributions of the latter five operators to neutral and charged current amplitudes are equivalent to a renormalization of the four operators $\mathcal{O}_{BW}$, $\mathcal{O}_{DW}$, $\mathcal{O}_{DB}$, and $\mathcal{O}_{\Phi,1}$ which contribute at tree level. Although the number of parameters of this model is apparently rather large, it was shown in Ref. [15] that in the Z -peak subtracted representation only two parameters remain in the process $e^+e^- \rightarrow \bar{f}f$, since f_{BW}^r and $f_{\Phi,1}^r$ are fully reabsorbed in the input parameters Γ_Z^u and $s_W^2(M_Z^2)$.

In the Z -peak subtracted representation, the observables $\sigma_{\mu\mu}$ and σ_{FB} are given by [15]:

$$\sigma_{\mu\mu}(s) = \sigma_{\mu\mu}^{(\gamma)}(s) + \sigma_{\mu\mu}^{(Z)}(s) + \sigma_{\mu\mu}^{(\gamma Z)}(s), \quad (28)$$

$$\sigma_{\mu\mu}^{(\gamma)}(s) = \tilde{\sigma}_{pt} \left[1 + 2\tilde{\Delta}\alpha(s) \right], \quad (29)$$

$$\sigma_{\mu\mu}^{(Z)}(s) = \tilde{\sigma}_{pt} |\chi_Z|^2 a^4 \left[1 - 2R(s) - 16s_W c_W \frac{\tilde{v}_l}{1 + \tilde{v}_l^2} V(s) \right], \quad (30)$$

$$\begin{aligned} \sigma_{\mu\mu}^{(\gamma Z)}(s) &= \tilde{\sigma}_{pt} 2\text{Re}\chi_Z a^2 \frac{\tilde{v}_l^2}{1 + \tilde{v}_l^2} \times \\ &\times \left[1 + \tilde{\Delta}\alpha(s) - R(s) - 8 \frac{s_W c_W}{\tilde{v}_l} V(s) \right], \end{aligned} \quad (31)$$

and

$$\sigma_{\text{FB}}(s) = \sigma_{\text{FB}}^{(Z)}(s) + \sigma_{\text{FB}}^{(\gamma Z)}(s), \quad (32)$$

$$\sigma_{\text{FB}}^{(Z)}(s) = \frac{3}{4} \tilde{\sigma}_{pt} |\chi_Z|^2 a^4 \frac{4\tilde{v}_l^2}{(1 + \tilde{v}_l^2)^2} \left[1 - 2R(s) - 8 \frac{s_W c_W}{\tilde{v}_l} V(s) \right], \quad (33)$$

$$\sigma_{\text{FB}}^{(\gamma Z)}(s) = \frac{3}{4} \tilde{\sigma}_{pt} 2\text{Re}\chi_Z a^2 \frac{1}{1 + \tilde{v}_l^2} \left[1 + \tilde{\Delta}\alpha(s) - R(s) \right]. \quad (34)$$

Here: $\tilde{\sigma}_{pt} = \frac{4\pi\alpha^2}{3s}$, $a^2 = \frac{3\Gamma_Z^l}{\alpha M_Z}$, $\alpha \equiv \alpha(0) \simeq 1/137$, $s_W^2 \equiv \sin^2 \theta_W^{eff}$ as in (19), and $\tilde{v} = 1 - 4s_W^2$.

Contributions of anomalous gauge couplings (AGC) to the above equations can be written as:

$$\tilde{\Delta}^{(AGC)}\alpha(s) = -8\pi\alpha \frac{s}{\Lambda^2} [f_{DW}^r + f_{DB}^r], \quad (35)$$

$$R^{(AGC)}(s) = \frac{8\pi\alpha}{s_W^2 c_W^2} \frac{(s - M_Z^2)}{\Lambda^2} [c_W^4 f_{DW}^r + s_W^4 f_{DB}^r], \quad (36)$$

$$V^{(AGC)}(s) = \frac{8\pi\alpha}{s_W c_W} \frac{(s - M_Z^2)}{\Lambda^2} [c_W^2 f_{DW}^r - s_W^2 f_{DB}^r]. \quad (37)$$

In these equations, Λ is the mass scale of the new interaction, and f_{DW}^r and f_{DB}^r are the renormalized couplings that are associated to the six-dimensional effective operators

$$\mathcal{O}_{DW} = \text{Tr} \left([D_\mu, \hat{W}_{\nu\rho}] [D^\mu, \hat{W}^{\nu\rho}] \right), \quad (38)$$

$$\mathcal{O}_{DB} = -\frac{g'^2}{2} (\partial_\mu B_{\nu\rho}) (\partial^\nu B^{\nu\rho}), \quad (39)$$

that are involved in the effective Lagrangian [4] and survive in the Z -peak subtracted representation.

It is convenient to express the deviations of the cross section and of the forward-backward cross section from the SM prediction, at energies away from the Z peak, in the approximation based on the simplified assumption *ii*), and neglecting numerically irrelevant small contributions. The result is:

$$\Delta\sigma_{\mu\mu} \approx \sigma_{pt} \left(-8\pi\alpha \frac{s}{\Lambda^2} \right) \left[(2.4 f_{DW}^r + 1.8 f_{DB}^r) + a^2 \text{Re}\chi_Z (2.4 f_{DW}^r + 0.2 f_{DB}^r) \right] \quad (40)$$

and

$$\Delta\sigma_{\text{FB}} \approx \frac{3}{4} \sigma_{pt} \left(-8\pi\alpha \frac{s}{\Lambda^2} \right) \left[(2.4 f_{DW}^r + 0.2 f_{DB}^r) + a^2 \text{Re}\chi_Z (2.4 f_{DW}^r + 1.8 f_{DB}^r) \right]. \quad (41)$$

The corresponding expressions for the new observables σ_+ and σ_- can be easily derived from the relation (18) and the formulae for $\sigma_{\mu\mu}$ and σ_{FB} . The result is:

$$\Delta\sigma_+ \approx \sigma_{pt} \left(-8\pi\alpha \frac{s}{\Lambda^2} \right) (1 + a^2 \text{Re}\chi_Z) (2.4 f_{DW}^r + 1.0 f_{DB}^r), \quad (42)$$

and

$$\Delta\sigma_- \approx \sigma_{pt} \left(-8\pi\alpha \frac{s}{\Lambda^2} \right) (1 - a^2 \text{Re}\chi_Z) 0.8 f_{DB}^r. \quad (43)$$

Fig. 3a shows the relative deviation $\Delta\sigma_+/\sigma_+^{SM}$ induced by the anomalous gauge couplings with $(f_{DW}^r, f_{DB}^r) = (\pm 1, \pm 1)$ and $\Lambda = 1$ TeV, as numerical examples. In contrast to the Z' case (Fig. 1a), the sign of the deviation due to anomalous gauge couplings is not defined, i.e., it may be either positive or negative at energies up to LEP2. However, as Eq. (42) shows, characteristic energy correlations of the deviation occur also in this case. Indeed, for certain values of the parameters $(f_{DW}^r, f_{DB}^r = (1, 1); (1, -1))$ the energy behaviour of the deviation due to anomalous gauge couplings is the same as that in Fig. 1a for the Z' , while for other choices of the parameters $(f_{DW}^r, f_{DB}^r = (-1, 1); (-1, -1))$ the behaviour is opposite to the Z' case.

Fig. 3b shows the energy behaviour of $\Delta\sigma_-/\sigma_-^{SM}$ for the same choice of the parameters. Such behaviour is qualitatively similar to the Z' case, in particular regarding the energy correlations of the sign of the deviation. Moreover, from Eq. (43), $\Delta\sigma_-$ depends on the couplings f_{DB}^r only, and therefore just a one-parameter fit to current (and expected) experimental data is required for this parameter in our analysis.

Concerning the distinction of the two potential sources of nonstandard effects considered here, Z' vs anomalous gauge couplings, one can use the negative sign property at LEP2 of $\Delta\sigma_+$ as well as $\Delta\sigma_{\mu\mu}$ and $\Delta\sigma_{\text{FB}}$, which holds for the former case (Table 1) but not for the latter one. The hatched area in the (f_{DW}^r, f_{DB}^r) plane of Fig. 4 corresponds to the values of anomalous gauge couplings such that the deviations of Eqs. (40)-(42) are negative at LEP2. This is a ‘confusion’ region, where both nonstandard mechanisms can induce the same deviations from the SM prediction, and therefore cannot be distinguished through the property of negative sign deviation

of the observables introduced here. On the contrary, the white area on the left of the hatched one is determined by values of the anomalous couplings such that the corresponding (positive) deviations are not allowed to the Z' and, in this sense, the source of the nonstandard effect can be distinguished.

Another condition can be derived by considering that Eqs. (24) and (25) imply, for the Z' , that

$$\frac{\Delta\sigma_-}{\Delta\sigma_+} = \frac{1 - a_l^2 \text{Re}\chi_Z}{1 + a_l^2 \text{Re}\chi_Z} \cdot \frac{v_l'^2 - a_l'^2}{v_l'^2 + a_l'^2}, \quad (44)$$

with, of course, $|(v_l'^2 - a_l'^2)/(v_l'^2 + a_l'^2)| \leq 1$. For, e.g., $\sqrt{s} = 190$ GeV, the above ratio is limited by

$$\left| \frac{\Delta\sigma_-}{\Delta\sigma_+} \right| \leq 0.36. \quad (45)$$

Taking into account Eqs. (42) and (43) for the case of the anomalous gauge couplings:

$$\frac{\Delta\sigma_-}{\Delta\sigma_+} = \frac{1 - a_l^2 \text{Re}\chi_Z}{1 + a_l^2 \text{Re}\chi_Z} \cdot \frac{0.8f_{DB}^r}{2.4f_{DW}^r + 1.0f_{DB}^r}, \quad (46)$$

and, comparing with the numerical bound (45), one finds that the actual ‘confusion’ region between the two kinds of mechanism is limited by the lines 4 and 5 of Fig. 4, and therefore is somewhat reduced.

Finally, the contours in Fig. 5 represent the upper bounds on f_{DW}^r and f_{DB}^r allowed by the combination of σ_+ and σ_- at the 2- σ level, in the case of no deviation observed from the SM, using the current LEP2 data of Table 2 and the expected experimental uncertainty with the future full luminosity. Numerical values of the anomalous gauge couplings included in the area between the two ellipses give rise to deviations from the SM that are, potentially, still in the reach of future measurements at LEP2. The values of the anomalous couplings within the shaded area produce deviations which can also originate from the Z' , and in this sense identify the so-called ‘confusion’ area.

Acknowledgements

AAB and AAP gratefully acknowledge the support of the University of Trieste.

References

- [1] For a review see, e.g., L. Hewett and T. G. Rizzo, Phys. Rep. **C183** (1989) 193.
- [2] J.A. Valls, report FERMILAB-Conf-97/135-E.
- [3] Physics at LEP2, Proceedings of the Workshop, Geneva, Switzerland (1996), CERN 96-01, G. Altarelli, T. Sjöstrand and F. Zwirner, eds.

- [4] See, e.g., K. Hagiwara, S. Ishihara, R. Szalapski and D. Zeppenfeld, Phys. Lett. **B283** (1992) 353; Phys. Rev. D **48** (1993) 2182.
- [5] M. Consoli, W. Hollik and F. Jegerlehner, *in* Z physics at LEP1, ed. G. Altarelli, R. Kleiss and C. Verzegnassi, vol.1, p.7, 1989.
- [6] G. Altarelli, R. Casalbuoni, D. Dominici, F. Feruglio and R. Gatto, Nucl. Phys. **B342** (1990) 15.
- [7] G. Altarelli, B. Mele and M. Ruiz Altaba, Z. Phys. C **45** (1989) 109; (E) C **47** (1990) 676.
- [8] A. Leike, Z. Phys. C **62** (1994) 265.
- [9] P. Osland and A.A. Pankov, Phys. Lett. **B403** (1997) 93; **B406** (1997) 328.
- [10] L3 collaboration, M. Acciarri et al., preprint CERN-PPE/97-52, 1997.
- [11] OPAL Collaboration, K. Ackerstaff et al., preprint CERN-PPE/97-101, 1997.
- [12] ALEPH Collaboration, Contribution to the 1997 EPS-HEP Conference, Jerusalem, August 19-26, 1997, Ref.# 602.
- [13] DELPHI collaboration, A. Behrmann et al., preprint DELPHI 97-132 CONF 110, 1997.
- [14] S. Riemann, FORTRAN program ZEFIT Version 4.2;
D. Bardin et al., preprint CERN-TH. 6443/92, CERN, 1992.
- [15] F. M. Renard and C. Verzegnassi, Phys. Rev. D **52** (1995) 1369; D **53** (1996) 1290.

Figure captions

Fig. 1a Relative deviation $\Delta\sigma_+/\sigma_+^{SM}$ induced by a ‘sequential’ Z' with $v'_l = v_l$ and $a'_l = a_l$. $M_{Z'} = 500$ GeV (line 1) and $M_{Z'} = 700$ GeV (line 2). Notice $\Delta\sigma_+/\sigma_+^{SM} = 0$ at $\sqrt{s} = M_Z/\sqrt{1+a_l^2} \simeq 78$ GeV and $\sqrt{s} = M_Z$.

Fig. 1b Relative deviation $\Delta\sigma_-/\sigma_-^{SM}$. Lines 1 and 2 correspond to $v_l'^2 - a_l'^2 = v_l^2 - a_l^2$, $M_{Z'} = 500$ GeV and $M_{Z'} = 700$ GeV, respectively. Lines 3 and 4 correspond to $v_l'^2 - a_l'^2 = -(v_l^2 - a_l^2)$ and $M_{Z'} = 500$ GeV, $M_{Z'} = 700$ GeV, respectively. Notice $\Delta\sigma_-/\sigma_-^{SM} = 0$ at $\sqrt{s} = M_Z$ and $\sqrt{s} = M_Z/\sqrt{1-a_l^2} \simeq 113$ GeV.

Fig. 2 Upper bounds on the Z' couplings (A_l , V_l) determined by $\sigma_\pm$ at the 2σ level. The solid contour is obtained from $e^+e^- \rightarrow \mu^+\mu^-$ with inputs summarized in Table 2. The dashed line is obtained with $L_{int} = 500$ pb $^{-1}$ and $\sqrt{s} = 190$ GeV.

Fig. 3a Relative deviation $\Delta\sigma_+/\sigma_+^{SM}$ induced by anomalous gauge couplings. Lines 1, 2, 3 and 4 correspond to: $(f_{DW}^r, f_{DB}^r)=(1, 1); (1, -1); (-1, 1); (-1, -1)$, respectively.

Fig. 3b Relative deviation of $\Delta\sigma_-/\sigma_-^{SM}$, parameters as in Fig. 3a.

Fig. 4 Straight lines in the two-dimensional plane (f_{DW}^r, f_{DB}^r) corresponding to $\Delta\sigma_{\mu\mu} = 0$ (line 1), $\Delta\sigma_+ = 0$ (line 2) and $\Delta\sigma_{FB} = 0$ (line 3). The hatched area is the ‘confusion’ area between Z' and anomalous gauge couplings effects when the deviations $\Delta\sigma_{\mu\mu}$, $\Delta\sigma_{FB}$ and $\Delta\sigma_+$ are negative at LEP2. The shaded area is the reduced ‘confusion’ area using also the solutions of (45) and (46) (lines 4 and 5).

Fig. 5 Upper bounds on the anomalous gauge couplings (f_{DW}^r, f_{DB}^r) determined by $\sigma_{\pm}$ at the 2σ level. Input parameters for solid and dotted lines are as in Fig. 2. The shaded area is the ‘confusion’ one, where Z' and anomalous gauge couplings can produce the same deviation.













